

EXCEPTIONAL ORTHOPAEDIC CARE BEGINS HERE

Ryan Livermore, M.D.

10100 East Shannon Woods Circle Suite 100 Wichita, KS 67226

(316) 219-8299

Rotator Cuff Repair

Weeks 1-4: Rest and Healing

Sling Immobilizer: At all times except exercises

HEP: Distal ROM with scapular retraction and posture

Manual scapular manipulation with patient lying on non-operative side

Supine passive FF in scapular plane to 120

Supine passive ER to 30

Weeks 4-6: Protective/Early Motion Phase

Sling Immobilizer: At all times except exercises

PROM: Forward flexion in scapular plane – No limits

External rotation 45 degrees Internal rotation 30 degrees

Therapeutic exercises: Codmans, wand exercises

Strengthening: RTC isometrics with arm in 0 deg abduction and neutral rotation

Scapular stabilization, no resistance Abdominal and trunk exercises

Weeks 7-12: Early Strengthening Phase

PROM/AAROM: FF/ER/IR - Full

Therapeutic exercises: -Cont wand exercises for ER/IR/FF

-Flexibility, horizontal adduction (post capsule stretching)

Strengthening: RTC isotonic strengthening exercises

- AROM: side-lying ER and supine FF in scapular plane
- Progress to standing FF
- ER/IR @ modified neutral w/ elastic bands

Progress to rhythmic stabilization exercises

Progress to closed chain exercises

Weeks 12+: Late Strengthening Phase

Progress isotonic strengthening: periscapular and RTC musculature

- Lat pull downs
- Row machine
- Chest press

Flexibility: side-lying post capsule stretch

Progress scapular stabilization program

Initiate isokinetic strengthening (IR/ER) in scapular plane

Begin light plyometrics at 16-18 weeks Individualize program to meet demands of sport specific requirements at 20-24 weeks Initiate throwing program for overhead athletes at 20-24 weeks