

EXCEPTIONAL ORTHOPAEDIC CARE BEGINS HERE

Ryan Livermore, M.D.

10100 East Shannon Woods Circle Suite 100 Wichita, KS 67226

(316) 219-8299

PHYSICAL THERAPY SHOULDER ARTHROSCOPY WITH COMBINED ANTERIOR, SUPERIOR, & POSTERIOR LABRAL REPAIR

Weeks 3-6: Phase I

<u>Sling Immobilizer:</u> At all times except for showering and exercise Exercises:

Passive ER to 45 and extension to neutral

Passive FF in scapular plane to 140

AROM wrist/elbow

Scapular "pinches"

Pain free submaximal deltoid isometrics

Modalities as needed

Advancement Criteria: ER to 45

FF in scapular plane to 140

Minimal pain and inflammation

Weeks 6-10: Phase II

Sling Immobilizer: At all times except for showering and exercise

Discontinue at week 8

Exercises:

Passive & Active assisted FF in scapular plane – no limits (wand exercises, pulleys)

Passive & Active assisted ER – no limits

Manual scapular side-lying stabilization exercises

IR/ER submaximal, pain free isometrics

Modalities as needed

Advancement Criteria:

FF to 160

ER to 60

Normal scapulohumeral rhythm

Minimal pain and inflamation

Weeks 10-14: Phase III

Exercises:

AAROM for full FF and ER

AAROM for IR – no limits

IR/ER/FF isotonic strengthening

Scapular and latissimus strengthening Humeral head stabilization exercises Begin biceps strengthing Progress IR/ER to 90/90 position if required General upper extremity flexibility exercises

Advancement Criteria:

Normal scapulohumeral rhythm Full upper extremity ROM Isokinetic IR/ER strength 85% of uninvolved side Minimal pain and inflamation

Weeks 14-18: Phase IV

Exercises:

Continue full upper extremity strengthening program
Continue upper extremity flexibility exercises
Activity-specific plyometrics program
Begin sport or activity related program
Address trunk and lower extremity demands

Discharge Criteria:

Isokinetic IR/ER strength equal to uninvolved side Independent HEP Independent, pain-free sport or activity specific program