

EXCEPTIONAL ORTHOPAEDIC CARE BEGINS HERE

Dr. Justin Strickland

10100 E. Shannon Woods Wichita, KS 67226 | 316-219-8299

Patella Dislocation/Subluxation Protocol (Conservative Management)

Attention: Treatment for first time dislocations with no loose body.

Days 0-6: NO THERAPY

- **Brace:** Wear knee immobilizer.
- Weight Bearing: NWB week 1. Can begin WBAT with crutches after 1 week.

Days 7-20: Begin PT

- **Brace:** Change to ROM knee brace at the end of the first week and set at 0-90 degrees.
- **ROM:** As tolerated.
- Strengthening: Focus on VMO and hip abduction strengthening.
- Weight Bearing: Begin WBAT with crutches after one week. Progress to full WB with no assistive device.

Days 21-41

- **Brace:** Unlock ROM brace.
- **ROM:** As tolerated.
- Strengthening: Focus on VMO and hip abduction strengthening.

Days 42+ (end of 6 weeks)

• **Brace:** DC ROM brace and put into Lateral J Brace.

Note: If the patella dislocates again then they will need MPFL reconstruction.

Patient progression during the time frames along with general modality and exercise choice is left up to the discretion of the treating therapist. If you have any questions regarding this protocol, please contact (316) 219-8299.